



Dust Processing in the M82 Superwind with JWST MIRI and NIRCам




Serena A. Cronin^{1,2}, Alberto D. Bolatto², Rebecca C. Levy³, Elizabeth Tarantino³, Martha L. Boyer³, Karl D. Gordon³
& the JWST Starbursts Collaboration


¹cronin@umd.edu, ²University of Maryland, College Park, ³Space Telescope Science Institute

Background & Data

- Galactic winds are critical to galaxy evolution. Winds may enrich the CGM, prevent accretion from the halo, and trigger star formation via re-accreting "cold fountains" [1].
- The formation of massive ($\gtrsim 10^4 M_\odot$), compact ($R \sim 1$ pc) star clusters and supernovae in starbursts can drive multiphase galactic winds [2].
- M82 is an archetypal starburst galaxy. At $d \sim 3.6$ Mpc away [3], we can study this wind in incredible detail!
- JWST (GO #1701; PI: A. Bolatto) has revealed a complex filamentary structure of Polycyclic Aromatic Hydrocarbons (PAHs) in the M82 wind [4, 5].
- PAHs are easily destroyed by hot phases, so they trace the cooler phase of the wind. *How does this cool material survive the gauntlet of a hot wind?*




PAH Ratios





- PAH emission is sensitive to grain size, temperature, and ionization [6]. Thus, PAH ratios can diagnose the physical state of the cool phase of the wind.
- The $11.3/7.7\mu\text{m}$ ratio increases with distance from the ionizing starburst, indicating PAHs tend to be larger and/or more neutral in the wind.
- $3.3\mu\text{m}$ emission is sensitive to the smallest grains. The distributions of $3.3/11.3\mu\text{m}$ and $3.3/7.7\mu\text{m}$ indicate that the smallest PAHs can survive out to at least ~ 2 kpc.

Wind Structure



- FilFinder locates filaments and extracts their properties, such as length, width, and orientation [7].
- (Left) **Skeletons and branches of filaments.** The M82 wind contains a complex network of filaments. Of the ~ 1000 features found, ~ 275 are considered significant (> 30 pc long).
- (Below, left) **Sizes of significant filaments.** Significant filaments tend to be $\sim 7\times$ long as they are wide.
- (Below, right) **Orientations of significant filaments.** Significant filaments tend to be oriented $\sim \pm 30^\circ$ from the minor axis ($0^\circ = \text{North}$). This is consistent with the opening angle of the molecular outflow cone (55° [8]).



- (Left) Wavelet power spectrum: $P_j \propto 2^{-j\gamma}$, where γ is the fractal dimension [9].
- Kelvin-Helmholtz instabilities create a thin, hot-cool mixing layer with $\gamma \sim 2.5$ [10].
- We apply a wavelet transform to regions of the $7.7\mu\text{m}$ image.
- Both sides of the wind have $\gamma \sim 2.5$, indicating the structure in both wind lobes arises from turbulence.

References

- [1] Veilleux+2020, A&ARv, 28, 2
- [2] Heckman+1990, ApJS, 74, 833
- [3] Dalcanton+2009, ApJS, 183, 67
- [4] Bolatto+2024, ApJ 967, 63
- [5] Fisher+2025, MNRAS, 538, 3068
- [6] Draine+2021, ApJ, 917, 3
- [7] Koch & Rosolowsky (2015), MNRAS, 452, 3435
- [8] Walter+2002, ApJL, 580, L21
- [9] Mayboroda & Spergel (2025), arXiv:2503.23202
- [10] Fielding+2020, ApJL, 894, L24

Support for JWST-GO-01701 is provided by NASA through a grant from the Space Telescope Science Institute.

Can't get enough of starburst-driven winds? Check out my recent paper on the warm, ionized phase of the NGC 253 galactic wind! arxiv.org/abs/2505.04707

